

D2 to M2 Procedure for D2-Brane DBI Effective Action

by J. Klusoň

*Department of Theoretical Physics and Astrophysics
Faculty of Science, Masaryk University
Kotlářská 2, 602 00, Brno
Czech Republic
E-mail: klu@physics.muni.cz*

ABSTRACT: We apply the procedure that was suggested in [arXiv:0806.1639] to the case of abelian D2-brane Dirac-Born-Infeld effective action and discuss its limitation. Then we suggest an alternative form of this procedure that is based on an existence of interpolating action proposed in [hep-th/9707113,hep-th/9707011].

KEYWORDS: D-branes, M-branes.

Contents

1. Introduction	1
2. D2-Brane Action	3
3. Gauged Theory for M2-brane	6
4. Conclusion	10
A. Direct Dimensional Reduction for Massive M2-brane	10

1. Introduction

It was proposed by Bagger and Lambert in collection of very nice papers [1, 2, 3] and independently by Gustavsson in [4] ¹ following earlier works [5, 6] that a certain class of $N = 8$ super-conformal theories in three dimensions are potential candidates for the world-volume description of multiple M2-branes in M-theory. These constructions are based on introducing of an algebraic structure known as Lie 3-algebra that is needed for closure of supersymmetry algebra. The metric versions of the above theories fall into two classes that depend on whether the invariant bilinear form in 3-algebra space is positive definite or indefinite. The original theories proposed by Bagger-Lambert are Euclidean theories with positive definite bilinear form while more recent proposals [24, 25] contain bilinear form that is indefinite and these Lie 3-algebras are known as Lorentzian 3-algebras.

It was claimed that the Lorentzian 3-algebra theories capture the low-energy world-volume dynamics of multiple parallel M2-branes. This model has the required classical symmetries, but has several unresolved problems. In particular, the classical theory has ghosts, X_{\pm} . Moreover, the ghost-free formulation seems directly equivalent to the non-conformal D2-brane theory as was argued in [39, 42, 46]. Explicitly, it was shown very clearly in [46] how it is possible-starting from $N = 8$ SYM- systematically and uniquely recovery the theory [39, 42].

Since the analysis presented in [46] is very nice and interesting it certainly deserves further study. In fact, since $2+1$ dim $N = 8$ SYM theory describes low-energy dynamics of N D2-branes one can ask the question whether it is possible to extend this analysis [46] ²

¹For related works, see [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 46, 47, 48, 49, 50, 51, 56, 57, 59, 61, 67, 68, 70, 75, 76, 77, 79, 80, 78, 81, 82, 83, 87, 89, 90, 91, 100, 101, 103]. For study of supergravity duals of these theories, see [43, 44, 45, 52, 53, 54, 55, 58, 60, 62, 63, 64, 65, 66, 69, 71, 72, 73, 74, 84, 85, 86, 88, 102].

²In what follows we call this analysis as EMP procedure.

when we take non-linear corrections into account. As the first step in this direction we try to apply EMP procedure to the case of single Dirac-Born-Infeld (DBI) action for D2-brane.

We start our analysis with the remarkable form of $2 + 1$ dimensional action that was proposed long ago in [92, 93]. This action is an interpolating action that-after appropriate integration of some world-volume fields-either describes D2-brane DBI effective action in massive Type IIA supergravity or the directly dimensional reduced gauged M2-brane action. We show that in linearized level this action is equivalent to the abelian form of the action given in [46] and hence can be considered as the starting point for non-linear generalization of EMP procedure. On the other hand we argue that naive application of EMP procedure in this action leads to a puzzle. Explicitly, we argue that there is a unique ground state of this new action with infinite coupling constant. This is different from we would expect since M2 to D2-reduction is based on a presumption that the vacuum expectation value of $\langle X_+ \rangle$ can take arbitrary constant value. Equivalently, we would expect an infinite number of ground states that differ by vacuum expectation values of X_+ .

In order to resolve this problem we suggest that the natural object for the definition of non-linear EMP procedure is gauged M2-brane action. More precisely, it is well known that in the case of IIA supergravity it is possible to introduce non-zero cosmological constant proportional to m^2 with m a mass parameter [94]. Such backgrounds are essential for the existence of D8-branes whose charge is proportional to m [95]. The action for massive 11-dimensional theory has the same content as the massless one ³

$$\hat{g}_{MN} , \quad \hat{C}_{MNK} , \quad M, N, K = 0, \dots, 11 . \quad (1.1)$$

The action for these fields is manifestly 11-dimensional Lorentz covariant but it does not correspond to a proper 11-dimensional theory because, in order to write down the action, we need to introduce an auxiliary non-dynamic vector field \hat{k}^M such that the Lie derivatives of the metric and 3-form potential with respect to it are zero:

$$\mathcal{L}_{\hat{k}} \hat{g}_{MN} = 0 , \quad \mathcal{L}_{\hat{k}} \hat{C}_{MNL} = 0 . \quad (1.2)$$

An existence of this Killing vector is crucial for definition of massive M2-brane. In fact, the world-volume theory of massive branes ⁴ was extensively studied in the past, for example [96, 97, 98, 99]. These actions have as a common characteristic that they are gauged sigma models. The gauged isometry is the same as an isometry that is needed in order to define the massive 11-dimensional supergravity theory. For example, the original un-gauged M2-brane action is the same object as in the massless theory, i.e. the corresponding massless M2-brane.

Let us again return to generalized EMP procedure. We argue that it can be naturally applied for gauged M2-brane action. As opposite to the original EMP procedure, where

³We use the following notation for the hats. Hats on target space fields indicate that they are 11-dimensional.

⁴These branes-that propagate in the background with non-zero cosmological constant-are called as "massive branes" as opposed to branes that propagate in a background with zero mass parameter. It is clear that all these branes are massive in the sense that their physical mass is nonzero.

the Yang-Mills coupling constant vector g_{YM}^I is replaced with a dynamical field X_+^I we replace the constant Killing vector \hat{k}^M with dynamical field \hat{X}_+^M . Then we can easily find manifestly covariant form of the generalized action with infinite number of ground states that differ by vacuum expectation values of \hat{X}_+^M .

It is remarkable that the gauged isometry that appears in massive M2-brane action is related to the gauge symmetry introduced in [46]. We hope that this observation will allow to find new geometrical interpretations of gauge symmetries that were introduced in [39, 42, 46].

The organization of this paper is as follows. In the next section (2) we introduce the interpolating D2-brane action and we argue that after appropriate redefinition of world-volume fields it agrees with the abelian version of D2-brane action introduced in [46]. In section (3) we apply EMP prescription for gauged M2-brane action and we find covariant and non-linear version of M2-brane action that has all desired properties. In conclusion (4) we outline our results and suggest possible extension of our work. Finally, in Appendix A we explicitly show that the dimensional reduction in gauged M2-brane action leads to the interpolating action introduced in section (2).

2. D2-Brane Action

We start with the action that was proposed in [92, 93]

$$\begin{aligned} S[X^m, X, V_\mu, B_\mu] = & \\ = -\tau_{M2} \int d^3\xi e^{-\Phi} \sqrt{-\det[g_{\mu\nu} + e^{2\Phi} F_\mu F_\nu]} + & \\ + \frac{\tau_{M2}}{3!} \int d^3\xi \epsilon^{\mu\nu\rho} [C_{\mu\nu\rho}^{(3)} + 6\pi\alpha' D_\mu X \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho] , & \end{aligned} \quad (2.1)$$

where

$$\begin{aligned} F_\mu &= D_\mu X + C_\mu^{(1)} , \\ D_\mu X &= \partial_\mu X + B_\mu , \\ \mathcal{F}_{\mu\nu} &= \partial_\mu V_\nu - \partial_\nu V_\mu - \frac{1}{2\pi\alpha'} b_{\mu\nu} \end{aligned} \quad (2.2)$$

and where

$$\begin{aligned} g_{\mu\nu} &= \partial_\mu X^m \partial_\nu X^n g_{mn} , \quad b_{\mu\nu} = b_{mn} \partial_\mu X^m \partial_\nu X^n , \\ C_{\mu\nu\rho}^{(3)} &= C_{mnk} \partial_\mu X^m \partial_\nu X^n \partial_\rho X^k , \quad C_\mu^{(1)} = C_m^{(1)} \partial_\mu X^m , \end{aligned} \quad (2.3)$$

where g_{mn}, b_{mn} are space-time metric and NS two form field respectively and where $C_{mnk}^{(3)}, C_m^{(1)}$ are Ramond-Ramond three and one forms respectively. Further, $X^m, m, n = 0, \dots, 9$ are

world-volume modes that describe embedding of D2-brane in the target space-time. Finally, τ_{M2} is D2-brane tension defined as $\tau_{M2} = \frac{1}{l_s^3}$.

The action (2.1) contains extra fields as opposite to usual DBI action for D2-brane. Firstly, if we integrate out B_μ we obtain

$$-\frac{e^\Phi \sqrt{-g} g^{\mu\nu} F_\nu}{\sqrt{1 + e^{2\Phi} g^{\mu\nu} F_\mu F_\nu}} + \pi \alpha' \epsilon^{\mu\nu\rho} \mathcal{F}_{\nu\rho} = 0 , \quad (2.4)$$

where we used the fact that

$$\sqrt{-\det[g_{\mu\nu} + e^{2\Phi} F_\mu F_\nu]} = \sqrt{-\det g} \sqrt{1 + e^{2\Phi} g^{\mu\nu} F_\mu F_\nu} . \quad (2.5)$$

Then if we insert (2.4) into (2.1) we obtain an action in the form

$$\begin{aligned} S = & -\tau_{M2} \int d^3 \xi e^{-\Phi} \sqrt{-\det[g_{\mu\nu} + 2\pi\alpha' \mathcal{F}_{\mu\nu}]} + \\ & + \frac{\tau_{M2}}{3!} \int d^3 \xi \epsilon^{\mu\nu\rho} (C_{\mu\nu\rho}^{(3)} - 6\pi\alpha' C_\mu^{(1)} \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho) \end{aligned} \quad (2.6)$$

that is standard form of D2-brane in massive Type IIA background and that reduces to the massless Type IIA background when $m = 0$.

In order to see that the action (2.6) is related to abelian reduction of the action given in [46] we take following background:

$$g_{mn} = \eta_{mn} , \quad \Phi = \Phi_0 = \text{const} , \quad C_m^{(1)} = C_{mnk}^{(3)} = 0 . \quad (2.7)$$

Further, let us impose static gauge

$$X^\mu = \xi^\mu , \quad \mu = 0, 1, 2 \quad (2.8)$$

so that

$$g_{\mu\nu} = \eta_{\mu\nu} + \delta_{ij} \partial_\mu X^i \partial_\nu X^j , \quad i, j = 3, \dots, 9 . \quad (2.9)$$

Then in the quadratic approximation the action (2.6) takes the form

$$\begin{aligned} S[X^m, X, V_\mu, B_\mu] = & -\tau_{M2} \int d^3 \xi e^{-\Phi_0} \sqrt{-\eta} - \tau_{M2} \int d^3 \xi \sqrt{-\eta} \left[\frac{1}{2} e^{-\Phi_0} \eta^{\mu\nu} \delta_{ij} \partial_\mu X^i \partial_\nu X^j + \right. \\ & \left. + \frac{e^{\Phi_0}}{2} \eta^{\mu\nu} F_\mu F_\nu \right] + \int d^3 \xi \epsilon^{\mu\nu\rho} (\pi\alpha' \tau_{M2}) D_\mu X \mathcal{F}_{\nu\rho} . \end{aligned} \quad (2.10)$$

As the next step we introduce the gauge theory coupling constant through the standard relations

$$e^{-\Phi_0} l_s^4 \tau_{M2} = \frac{1}{g_{YM}^2} , \quad \left(\tau_{M2} = \frac{1}{l_s^3} , \quad 2\pi\alpha' = l_s \right) \quad (2.11)$$

so that after rescaling

$$\sqrt{\tau_{M2}}e^{\Phi_0/2}X = \tilde{X} \ , \quad \sqrt{\tau_{M2}}e^{-\Phi_0/2}X^i = \tilde{X}^i \ , \quad \frac{1}{l_s^{5/2}}B_\mu = \tilde{B}_\mu \ , \quad l_s^{3/2}\mathcal{F}_{\mu\nu} = \tilde{\mathcal{F}}_{\mu\nu} \quad (2.12)$$

the action (2.10) takes the form

$$\begin{aligned} S[\tilde{X}^m, \tilde{X}, \tilde{V}_\mu, \tilde{B}] = & -\frac{1}{g_{YM}^2 l_s^2} \int d^3\xi \sqrt{-\eta} - \int d^3\xi \sqrt{-\eta} \left[\frac{1}{2} \eta^{\mu\nu} \delta_{ij} \partial_\mu \tilde{X}^i \partial_\nu \tilde{X}^j + \right. \\ & + \frac{1}{2} \eta^{\mu\nu} (\partial_\mu \tilde{X} + g_{YM} \tilde{B}_\mu) (\partial_\nu \tilde{X} + g_{YM} \tilde{B}_\nu) \Big] + \\ & + \int d^3\xi \epsilon^{\mu\nu\rho} \left[\frac{1}{2} \tilde{B}_\mu \tilde{\mathcal{F}}_{\nu\rho} + \frac{1}{2l_s^{3/2} g_{YM}} \partial_\mu \tilde{X} \tilde{\mathcal{F}}_{\nu\rho} \right] \end{aligned} \quad (2.13)$$

that has the same form as the abelian form of the action given in [46]⁵. Motivated by this result we perform the rescaling (2.12) in the action (2.6) and we obtain

$$\begin{aligned} S[\tilde{X}^m, \tilde{X}, \tilde{V}_\mu, \tilde{B}_\mu] = & - \int d^3\xi \sqrt{-\det \mathbf{A}_{\mu\nu}} + \int d^3\xi \epsilon^{\mu\nu\rho} \frac{1}{2} \tilde{B}_\mu \tilde{\mathcal{F}}_{\nu\rho} \ , \\ \mathbf{A}_{\mu\nu} = & \frac{1}{l_s^{8/3} g_{YM}^{4/3}} \eta_{\mu\nu} + g_{YM}^{2/3} l_s^{4/3} \partial_\mu \tilde{X}^i \partial_\nu \tilde{X}^j \delta_{ij} + \\ & + l_s^{4/3} g_{YM}^{2/3} (\partial_\mu \tilde{X} + g_{YM} \tilde{B}_\mu) (\partial_\nu \tilde{X} + g_{YM} \tilde{B}_\nu) \ , \end{aligned} \quad (2.14)$$

where we ignored term that contributes to the action as total derivative. Now we are ready to apply EMP procedure for (2.14). We introduce 8-dimensional vector g_{YM}^I as

$g_{YM}^I = (\overbrace{0, \dots, 0}^7, g_{YM})$, $I = 1, \dots, 8$, and "covariant derivative \tilde{D} "

$$\tilde{D}_\mu \tilde{X}^i = \partial_\mu \tilde{X}^i + g_{YM}^i \tilde{B}_\mu \ , \quad \tilde{D}_\mu \tilde{X} = \partial_\mu \tilde{X} + g_{YM} \tilde{B}_\mu \ . \quad (2.15)$$

Further, we rewrite g_{YM}^2 in manifest $SO(8)$ covariant manner as $g_{YM}^2 = g_{YM}^I g_{YM}^J \delta_{IJ} = |g_{YM}|^2$ and then we replace vector g_{YM}^I with dynamical field X_+^I so that the action (2.14) takes the form

$$\begin{aligned} S[\tilde{X}^I, \tilde{X}_+^I, \tilde{V}_\mu, \tilde{B}_\mu, C_I^\mu] = & \\ = & - \int d^3\xi \left(\sqrt{-\det \mathbf{A}_{\mu\nu}} + \epsilon^{\mu\nu\rho} \frac{1}{2} \tilde{B}_\mu \tilde{\mathcal{F}}_{\nu\rho} + C_I^\mu \partial_\mu \tilde{X}_+^I \right) \ , \\ \mathbf{A}_{\mu\nu} = & \frac{1}{l_s^{8/3} (X_+^I X_+^J \delta_{IJ})^{2/3}} \eta_{\mu\nu} + (X_+^I X_+^J \delta_{IJ})^{1/3} l_s^{4/3} \tilde{D}_\mu \tilde{X}^I \tilde{D}_\nu \tilde{X}^J \delta_{IJ} \ , \end{aligned} \quad (2.16)$$

where we introduced auxiliary field C_I^μ that renders \tilde{X}_+^I non-dynamical.

⁵This is true up to total derivative term since $\int d^3\xi \epsilon^{\mu\nu\rho} \partial_\mu \tilde{X} \tilde{\mathcal{F}}_{\nu\rho} = \int d^3\xi \partial_\mu [\epsilon^{\mu\nu\rho} \tilde{X} \tilde{\mathcal{F}}_{\nu\rho}] - \int d^3\xi \tilde{X} \partial_\mu (\epsilon^{\mu\nu\rho} \tilde{\mathcal{F}}_{\nu\rho}) = \int d^3\xi \partial(\dots)$.

Let us now analyze some properties of the action (2.16). We are mainly interested in the study of the ground state of this theory that has to solve the equations of motion that follow from the action (2.16). We presume that the ground state is characterized by following configuration of the world-volume fields

$$\tilde{X}_+^8 = v = \text{const} , \quad \tilde{D}_\mu \tilde{X}^I = 0 , \quad \tilde{B}_\mu = 0 , \quad \tilde{\mathcal{F}}_{\mu\nu} = 0 . \quad (2.17)$$

Firstly, the equation of motion for C_I^μ takes the form

$$\partial_\mu \tilde{X}_+^I = 0 \quad (2.18)$$

that is clearly obeyed by the ansatz (2.17). On the other hand the equation of motion for X_+^I takes the form

$$-\frac{1}{2} \frac{\delta \mathbf{A}_{\mu\nu}}{\delta X_+^I} (\mathbf{A}^{-1})^{\nu\mu} \sqrt{-\det \mathbf{A}} - \epsilon^{\mu\nu\rho} \frac{1}{2l_s^{1/2} (X_+^I X_+^J \delta_{IJ})^{3/2}} \partial_\mu \tilde{X} \tilde{\mathcal{F}}_{\nu\rho} = 0 . \quad (2.19)$$

Since for the ansatz (2.17) the matrix $\mathbf{A}_{\mu\nu}$ is equal to

$$\mathbf{A}_{\mu\nu} = \frac{1}{l_s^{8/3} v^{4/3}} \eta_{\mu\nu} \quad (2.20)$$

the equation (2.19) implies

$$\frac{1}{v^3} = 0 . \quad (2.21)$$

In other words the ground state corresponds to the point $v \rightarrow \infty$ that implies that there is unique ground state of the theory. As we argued in introduction this is not the same what we want since we would like to have a theory with infinite number of ground states that differ by vacuum expectation values of \tilde{X}_+ . In order to find solution of this problem we suggest an alternative procedure how to introduce X_+^I as a new dynamical variable. In the next section we present such an alternative procedure that is based on the fact that the action (2.6) can be considered as dimensional reduction of massive M2-brane.

3. Gauged Theory for M2-brane

Let us again consider the action (2.6) and determine the equations of motion for V_μ

$$\pi m \alpha' \epsilon^{\mu\nu\rho} (\partial_\nu V_\rho - \partial_\rho V_\mu) + \epsilon^{\mu\nu\rho} (\partial_\nu B_\rho - \partial_\rho B_\nu) = 0 . \quad (3.1)$$

Inserting (3.1) back to (2.6) we obtain the action in the form

$$\begin{aligned} S = & -\tau_{M2} \int d^3\xi \sqrt{-\det(e^{-2/3\Phi} g_{\mu\nu} + e^{4/3\Phi} F_\mu F_\nu)} + \\ & + \frac{\tau_{M2}}{6} \int d^3\xi \epsilon^{\mu\nu\rho} (C_{\mu\nu\rho}^{(3)} - 3D_\mu X B_{\nu\rho}^{(1)} + \frac{6}{m} B_\mu \partial_\nu B_\rho) . \end{aligned} \quad (3.2)$$

As was shown in [93] (and reviewed in appendix A) this action is very close to the action that one gets by direct dimensional reduction of the massive M2-brane that is also known as gauged M2-brane action. This action can be defined in the background with Killing vector isometry $\hat{k}^M(\hat{X})$. Then the gauged M2-brane action takes the form [93]

$$S = -\tau_{M2} \int d^3\xi \sqrt{-\det D_\mu \hat{X}^M D_\nu \hat{X}^N \hat{g}_{MN}} + \tau_{M2} \int d^3\xi \epsilon^{\mu\nu\rho} [D_\mu \hat{X}^M D_\nu \hat{X}^N D_\rho \hat{X}^K \hat{C}_{MNK} - \frac{6}{m} B_\mu \partial_\nu B_\rho] , \quad (3.3)$$

where the covariant derivative D_μ is defined as

$$D_\mu \hat{X}^M = \partial_\mu \hat{X}^M + B_\mu \hat{k}^M(\hat{X}) , \quad (3.4)$$

where B_μ is world-volume gauge field related to the Killing gauge isometry. To clarify meaning of this gauged form of the action let us consider following transformation

$$\delta_\eta \hat{X}^M(\xi) = \hat{X}'^M(\xi) - \hat{X}^M(\xi) = \eta(\xi) \hat{k}^M(\hat{X}) , \quad (3.5)$$

where $\eta(\xi)$ is a parameter of gauge transformations. This transformation immediately implies following transformation rules of background fields

$$\delta_\eta \hat{g}_{MN} = \eta \hat{k}^K \partial_K \hat{g}_{MN} , \quad \delta_\eta \hat{C}_{KMN} = \eta \hat{k}^L \partial_L \hat{C}_{KMN} , \quad \delta_\eta \hat{k}^K = \eta \hat{k}^L \partial_L \hat{k}^K \quad (3.6)$$

and transformation of covariant derivative

$$\delta_\eta D_\mu \hat{X}^M = \eta D_\mu \hat{X}^L \partial_L \hat{k}^M , \quad (3.7)$$

where we postulate following transformation rule for gauge field B_μ

$$\delta B_\mu = -\partial_\mu \eta . \quad (3.8)$$

Then

$$\delta(D_\mu \hat{X}^M D_\nu \hat{X}^N \hat{g}_{MN}) = \eta D_\mu \hat{X}^M (\partial_M \hat{k}^L \hat{g}_{LN} + \hat{g}_{ML} \partial_N \hat{k}^L + \partial_L \hat{g}_{MN}) D_\nu \hat{X}^N = 0 \quad (3.9)$$

since

$$\mathcal{L}_{\hat{k}} \hat{g}_{MN} = 0 . \quad (3.10)$$

In the same way we obtain that

$$\delta_\eta (D_\mu \hat{X}^M D_\nu \hat{X}^N D_\rho \hat{X}^K \hat{C}_{MNK}) = 0 , \quad \mathcal{L}_{\hat{k}} \hat{C}_{MNK} = 0 \quad (3.11)$$

hence we see that the action is invariant under transformations (3.5) and (3.6).

Having clarified the fact that the D2-brane action (2.6) is related to the gauged M2-brane action we now introduce modified EMP procedure to the action (3.3). As

the first step in our construction we will presume an existence of adapted system of coordinates where $\hat{k}^M = \text{const.}$ This is always possible to achieve in flat background $\hat{g}_{MN} = \eta_{MN}$, $\hat{C}_{MNK} = 0$. Further, in analogy with EMP prescription, we replace constant \hat{k}^M with dynamical field \hat{X}_+^M and add to the action term $\frac{1}{2}C_M^\mu \partial_\mu \hat{X}_+^M$ to render this field non-dynamical.

Further we rewrite the Wess-Zumino term in (3.3) as

$$\pi\alpha' \epsilon^{\mu\nu\rho} B_\mu \mathcal{F}_{\nu\rho} + (\pi\alpha')^2 m \epsilon^{\mu\nu\rho} V_\mu \partial_\nu V_\rho . \quad (3.12)$$

In fact it is easy to see that now the equation of motion for V_μ that follow from (3.12) implies

$$\frac{1}{m\pi\alpha'} (\partial_\mu B_\nu - \partial_\nu B_\mu) = -(\partial_\mu V_\nu - \partial_\nu V_\mu) \quad (3.13)$$

and hence when we insert it back to (3.12) we obtain the last term in (3.3). Note also that this expression is invariant under η transformations (up to total derivative) since

$$\delta(\epsilon^{\mu\nu\rho} B_\mu \mathcal{F}_{\nu\rho}) = \epsilon^{\mu\nu\rho} \partial_\mu \eta \mathcal{F}_{\nu\rho} = -\eta \epsilon^{\mu\nu\rho} \partial_\mu \partial_\nu A_\rho + \eta \epsilon^{\mu\nu\rho} \partial_\mu \partial_\rho A_\nu = 0 . \quad (3.14)$$

In summary we derive the action in the form

$$\begin{aligned} S = & -\tau_{M2} \int d^3\xi [\sqrt{-\det \mathcal{G}_{\mu\nu}} + \frac{1}{2} \sqrt{-\det \mathcal{G}_{\mu\nu}} C_\nu^N \eta_{NM} (\mathcal{G}^{-1})^{\nu\mu} \partial_\mu \hat{X}_+^M + \\ & + \frac{\tau_{M2}}{3!} \int d^3\xi \epsilon^{\mu\nu\rho} [6\pi\alpha' B_\mu \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho] , \end{aligned} \quad (3.15)$$

where we added term $\frac{1}{2}C_\mu^M (\mathcal{G}^{-1})^{\mu\nu} \eta_{MN} \partial_\nu \hat{X}_+^N$ that renders \hat{X}_+^M constant on-shell and where we also introduced "generalized metric" $\mathcal{G}_{\mu\nu}$

$$\mathcal{G}_{\mu\nu} = (\partial_\mu \hat{X}^M + B_\mu \hat{X}_+^M) \eta_{MN} (\partial_\nu \hat{X}^N + B_\nu \hat{X}_+^N) . \quad (3.16)$$

Following [46] we introduce field \hat{X}_-^M and add to the action an expression

$$\frac{1}{2} \sqrt{-\det \mathcal{G}} \partial_\mu \hat{X}_-^M \eta_{MN} (\mathcal{G}^{-1})^{\mu\nu} \partial_\nu \hat{X}_+^N$$

in order the action will be invariant under additional shift symmetry

$$\delta C_\mu^M = \partial_\mu \lambda^M , \quad \delta \hat{X}_-^M = \lambda^M . \quad (3.17)$$

Then the final form of the action takes the form

$$\begin{aligned} S = & -\tau_{M2} \int d^3\xi \sqrt{-\det \mathcal{G}_{\mu\nu}} (1 + \\ & + \frac{1}{2} (C_\mu^M - \partial_\mu \hat{X}_-^M) \eta_{MN} (\mathcal{G}^{-1})^{\mu\nu} \partial_\nu \hat{X}_+^M) + \\ & + \frac{\tau_{M2}}{3!} \int d^3\xi \epsilon^{\mu\nu\rho} [6\pi\alpha' B_\mu \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho] \end{aligned} \quad (3.18)$$

Note also that in order to achieve that \hat{X}_+^M is constant on-shell and that the action possesses additional shift symmetry we can consider more general form of the action

$$S = -\tau_{M2} \int d^3\xi \sqrt{-\det \mathcal{G}_{\mu\nu}} \sqrt{1 + (C_\mu^M - \partial_\mu \hat{X}_-^M) \eta_{MN} (\mathcal{G}^{-1})^{\mu\nu} \partial_\nu \hat{X}_+^N} - \frac{\tau_{M2}}{3!} \int d^3\xi \epsilon^{\mu\nu\rho} [6\pi\alpha' B_\mu \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho] \quad (3.19)$$

that can be finally written in a suggestive form as

$$S = -\tau_{M2} \int d^3\xi \sqrt{-\det \mathbf{A}_{\mu\nu}} - \frac{\tau_{M2}}{3!} \int d^3\xi \epsilon^{\mu\nu\rho} [6\pi\alpha' B_\mu \mathcal{F}_{\nu\rho} + 6m(\pi\alpha')^2 V_\mu \partial_\nu V_\rho] \quad \mathbf{A}_{\mu\nu} = \mathcal{G}_{\mu\nu} + (C_\mu^M - \partial_\mu \hat{X}_-^M) \eta_{MN} \partial_\nu \hat{X}_+^N . \quad (3.20)$$

Let us now study properties of the action (3.20). Clearly it is invariant under shift symmetry (3.17). Further, the variation of this action with respect to C_μ^M implies

$$\eta_{NM} \partial_\nu \hat{X}_+^M (\mathbf{A}^{-1})^{\nu\mu} \sqrt{-\det \mathbf{A}} = 0 \quad (3.21)$$

that implies $\partial_\nu \hat{X}_+^M = 0$. Let us again presume the ground state of the theory in the form

$$B_\mu = \hat{X}_-^M = C_\mu^M = V_\mu = 0 , \quad \hat{X}_+^M = v^M . \quad (3.22)$$

It is easy to see that the equations of motion for \hat{X}^M , B_μ , C_μ^M and \hat{X}_-^M are obeyed for this ansatz. Finally, the problematic equation of motion for \hat{X}_+^M takes the form

$$B_\mu \eta_{MN} (\partial_\nu \hat{X}_+^N + B_\nu \hat{X}_+^N) (\mathbf{A}^{-1})^{\nu\mu} \sqrt{-\det \mathbf{A}} + \frac{1}{2} \partial_\mu [\eta_{MN} \partial_\nu \hat{X}_-^N (\mathbf{A}^{-1})^{\nu\mu} \sqrt{-\det \mathbf{A}}] = 0 \quad (3.23)$$

that is clearly solved by (3.22) for any v^M . Finally, let us impose the static gauge in the following form

$$\hat{X}^\mu = \xi^\mu , \quad \mu, \nu = 0, 1, 2 \\ C_\mu^\nu = 0 , \quad \hat{X}_+^\mu = \hat{X}_-^\mu = 0 \quad (3.24)$$

so that the matrix $\mathbf{A}_{\mu\nu}$ takes the form

$$\mathbf{A}_{\mu\nu} = \eta_{\mu\nu} + (\partial_\mu \hat{X}^I + B_\mu \hat{X}_+^I) (\partial_\nu \hat{X}^J + B_\nu \hat{X}_+^J) \delta_{IJ} + (C_\mu^I - \partial_\mu \hat{X}_-^I) \delta_{IJ} \partial_\nu \hat{X}_+^J , \quad I, J = 1, \dots, 8 . \quad (3.25)$$

Then the action up to quadratic approximation can be written as

$$S = -\tau_{M2} \int d^3\xi \sqrt{-\eta} - \tau_{M2} \int d^3\xi \sqrt{-\det \eta} \mathcal{L} , \quad (3.26)$$

where the Lagrangian density takes the form

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \eta^{\mu\nu} (\partial_\mu \hat{X}^I + B_\mu \hat{X}_+^I) \delta_{IJ} (\partial_\nu \hat{X}^J + B_\nu \hat{X}_+^J) \delta_{IJ} + \\ & + \frac{1}{2} \eta^{\mu\nu} (C_\mu^I - \partial_\mu \hat{X}_-^I) \partial_\nu \hat{X}_+^J \delta_{IJ} - \\ & - \epsilon^{\mu\nu\rho} (\pi \alpha' B_\mu \mathcal{F}_{\nu\rho} + m (\pi \alpha')^2 V_\mu \partial_\nu V_\rho) . \end{aligned} \quad (3.27)$$

that is again very close to the abelian form of the action given in [46] and provides further support of our construction.

4. Conclusion

Let us summarize our results. We studied EMP procedure for Dirac-Born-Infeld action for D2-brane and we found its limitation. Then we suggested an alternative form of this procedure that is based on a formulation of gauged M2-brane action. This fact however implies that the theory should be defined in background with non-zero mass parameter m and this observation certainly deserves better understanding and more detailed study. Further, it would be also interesting to develop BRST Hamiltonian treatment of the action (3.20) and compare it with the similar analysis that was given in [24]. Finally, it will be extremely interesting to see whether there exists an non-abelian extension of the action (3.20). We hope to return to these problems in future.

Acknowledgement

This work was supported by the Czech Ministry of Education under Contract No. MSM 0021622409.

A. Direct Dimensional Reduction for Massive M2-brane

In this appendix we show that the gauged M2-brane action upon direct dimensional reduction in the direction X associated to the gauged isometry reduces to the action (3.2). To begin with we choose coordinates that are adapted to the isometry so that $\hat{k}^M = \delta^{Mx}$ and we split eleven coordinates \hat{X}^M into the ten 10-dimensional $X^m, m = 0, \dots, 9$ and the extra scalar $\hat{X}^x \equiv X$. Using the relations between the 11-dimensional and 10-dimensional fields

$$\begin{aligned} \hat{g}_{xx} &= e^{\frac{4}{3}\Phi} , \quad \hat{g}_{mx} = e^{\frac{4}{3}\Phi} C_m^{(1)} , \\ \hat{g}_{mn} &= e^{-\frac{2}{3}\Phi} g_{mn} + e^{\frac{4}{3}\Phi} C_m^{(1)} C_n^{(1)} , \\ \hat{C}_{mnk} &= C_{mnk}^{(3)} , \quad \hat{C}_{mnx} = B_{mn} \end{aligned} \quad (A.1)$$

it is straightforward to see that

$$\mathbf{A}_{\mu\nu} = e^{-\frac{2}{3}\Phi} \partial_\mu X^m \partial_\nu X^n g_{mn} + e^{\frac{4}{3}\Phi} (\partial_\mu X + B_\mu + C_\mu^{(1)}) (\partial_\nu X + B_\nu + C_\nu^{(1)}) . \quad (\text{A.2})$$

Then if we insert (A.2) together with (A.1) into the action (3.3) we easily obtain that it reduces to the action (3.2).

References

- [1] J. Bagger and N. Lambert, “*Modeling multiple M2’s*,” Phys. Rev. D **75** (2007) 045020 [arXiv:hep-th/0611108].
- [2] J. Bagger and N. Lambert, “*Gauge Symmetry and Supersymmetry of Multiple M2-Branes*,” Phys. Rev. D **77** (2008) 065008 [arXiv:0711.0955 [hep-th]].
- [3] J. Bagger and N. Lambert, “*Comments On Multiple M2-branes*,” JHEP **0802** (2008) 105 [arXiv:0712.3738 [hep-th]].
- [4] A. Gustavsson, “*Selfdual strings and loop space Nahm equations*,” JHEP **0804** (2008) 083 [arXiv:0802.3456 [hep-th]].
- [5] A. Basu and J. A. Harvey, “*The M2-M5 brane system and a generalized Nahm’s equation*,” Nucl. Phys. B **713** (2005) 136 [arXiv:hep-th/0412310].
- [6] J. H. Schwarz, “*Superconformal Chern-Simons theories*,” JHEP **0411** (2004) 078 [arXiv:hep-th/0411077].
- [7] S. Mukhi and C. Papageorgakis, “*M2 to D2*,” JHEP **0805** (2008) 085 [arXiv:0803.3218 [hep-th]].
- [8] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, “*N = 8 Superconformal Chern-Simons Theories*,” JHEP **0805** (2008) 025 [arXiv:0803.3242 [hep-th]].
- [9] D. S. Berman, L. C. Tadrowski and D. C. Thompson, “*Aspects of Multiple Membranes*,” Nucl. Phys. B **802** (2008) 106 [arXiv:0803.3611 [hep-th]].
- [10] M. Van Raamsdonk, “*Comments on the Bagger-Lambert theory and multiple M2-branes*,” JHEP **0805** (2008) 105 [arXiv:0803.3803 [hep-th]].
- [11] A. Morozov, “*On the Problem of Multiple M2 Branes*,” JHEP **0805** (2008) 076 [arXiv:0804.0913 [hep-th]].
- [12] N. Lambert and D. Tong, “*Membranes on an Orbifold*,” arXiv:0804.1114 [hep-th].
- [13] J. Distler, S. Mukhi, C. Papageorgakis and M. Van Raamsdonk, “*M2-branes on M-folds*,” JHEP **0805** (2008) 038 [arXiv:0804.1256 [hep-th]].
- [14] U. Gran, B. E. W. Nilsson and C. Petersson, “*On relating multiple M2 and D2-branes*,” arXiv:0804.1784 [hep-th].
- [15] P. M. Ho, R. C. Hou and Y. Matsuo, “*Lie 3-Algebra and Multiple M2-branes*,” JHEP **0806** (2008) 020 [arXiv:0804.2110 [hep-th]].
- [16] J. Gomis, A. J. Salim and F. Passerini, “*Matrix Theory of Type IIB Plane Wave from Membranes*,” arXiv:0804.2186 [hep-th].
- [17] E. A. Bergshoeff, M. de Roo and O. Hohm, “*Multiple M2-branes and the Embedding Tensor*,” Class. Quant. Grav. **25** (2008) 142001 [arXiv:0804.2201 [hep-th]].

- [18] K. Hosomichi, K. M. Lee and S. Lee, “*Mass-Deformed Bagger-Lambert Theory and its BPS Objects*,” arXiv:0804.2519 [hep-th].
- [19] G. Papadopoulos, “*M2-branes, 3-Lie Algebras and Plucker relations*,” JHEP **0805** (2008) 054 [arXiv:0804.2662 [hep-th]].
- [20] J. P. Gauntlett and J. B. Gutowski, “*Constraining Maximally Supersymmetric Membrane Actions*,” arXiv:0804.3078 [hep-th].
- [21] G. Papadopoulos, “*On the structure of k -Lie algebras*,” Class. Quant. Grav. **25** (2008) 142002 [arXiv:0804.3567 [hep-th]].
- [22] P. M. Ho and Y. Matsuo, “*M5 from M2*,” JHEP **0806** (2008) 105 [arXiv:0804.3629 [hep-th]].
- [23] J. Gomis, G. Milanesi and J. G. Russo, “*Bagger-Lambert Theory for General Lie Algebras*,” JHEP **0806** (2008) 075 [arXiv:0805.1012 [hep-th]].
- [24] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni and H. Verlinde, “ *$N=8$ superconformal gauge theories and M2 branes*,” arXiv:0805.1087 [hep-th].
- [25] P. M. Ho, Y. Imamura and Y. Matsuo, “*M2 to D2 revisited*,” JHEP **0807** (2008) 003 [arXiv:0805.1202 [hep-th]].
- [26] A. Morozov, “*From Simplified BLG Action to the First-Quantized M-Theory*,” arXiv:0805.1703 [hep-th].
- [27] Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, “*Janus field theories from multiple M2 branes*,” arXiv:0805.1895 [hep-th].
- [28] H. Fuji, S. Terashima and M. Yamazaki, “*A New $N=4$ Membrane Action via Orbifold*,” arXiv:0805.1997 [hep-th].
- [29] P. M. Ho, Y. Imamura, Y. Matsuo and S. Shiba, “*M5-brane in three-form flux and multiple M2-branes*,” arXiv:0805.2898 [hep-th].
- [30] Y. Song, “*Mass Deformation of the Multiple M2 Branes Theory*,” arXiv:0805.3193 [hep-th].
- [31] I. Jeon, J. Kim, N. Kim, S. W. Kim and J. H. Park, “*Classification of the BPS states in Bagger-Lambert Theory*,” arXiv:0805.3236 [hep-th].
- [32] M. Li and T. Wang, “*M2-branes Coupled to Antisymmetric Fluxes*,” arXiv:0805.3427 [hep-th].
- [33] K. Hosomichi, K. M. Lee, S. Lee, S. Lee and J. Park, “ *$N=4$ Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets*,” arXiv:0805.3662 [hep-th].
- [34] S. Banerjee and A. Sen, “*Interpreting the M2-brane Action*,” arXiv:0805.3930 [hep-th].
- [35] H. Lin, “*Kac-Moody Extensions of 3-Algebras and M2-branes*,” arXiv:0805.4003 [hep-th].
- [36] P. De Medeiros, J. Figueroa-O’Farril and E. Mendez-Escobar, “*Lorentzian Lie 3-algebras and their Bagger-Lambert moduli space*,” arXiv:0805.4363 [hep-th].
- [37] A. Gustavsson, “*One-loop corrections to Bagger-Lambert theory*,” arXiv:0805.4443 [hep-th].
- [38] J. Figueroa-O’Farrill, “*Lorentzian Lie n -algebras*,” arXiv:0805.4760 [math.RT].
- [39] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, “*Ghost-Free Superconformal Action for Multiple M2-Branes*,” arXiv:0806.0054 [hep-th].

- [40] J. H. Park and C. Sochichiu, “*Single M5 to multiple M2: taking off the square root of Nambu-Goto action,*” arXiv:0806.0335 [hep-th].
- [41] F. Passerini, “*M2-Brane Superalgebra from Bagger-Lambert Theory,*” arXiv:0806.0363 [hep-th].
- [42] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, “*Supersymmetric Yang-Mills Theory From Lorentzian Three-Algebras,*” arXiv:0806.0738 [hep-th].
- [43] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, “*N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals,*” arXiv:0806.1218 [hep-th].
- [44] C. Ahn, “*Holographic Supergravity Dual to Three Dimensional N=2 Gauge Theory,*” arXiv:0806.1420 [hep-th].
- [45] M. Benna, I. Klebanov, T. Klose and M. Smedback, “*Superconformal Chern-Simons Theories and AdS₄/CFT₃ Correspondence,*” arXiv:0806.1519 [hep-th].
- [46] B. Ezhuthachan, S. Mukhi and C. Papageorgakis, “*D2 to D2,*” JHEP **0807** (2008) 041 [arXiv:0806.1639 [hep-th]].
- [47] S. Cecotti and A. Sen, “*Coulomb Branch of the Lorentzian Three Algebra Theory,*” arXiv:0806.1990 [hep-th].
- [48] E. A. Bergshoeff, M. de Roo, O. Hohm and D. Roest, “*Multiple Membranes from Gauged Supergravity,*” arXiv:0806.2584 [hep-th].
- [49] P. de Medeiros, J. Figueroa-O’Farrill and E. Mendez-Escobar, “*Metric Lie 3-algebras in Bagger-Lambert theory,*” arXiv:0806.3242 [hep-th].
- [50] M. Blau and M. O’Loughlin, “*Multiple M2-Branes and Plane Waves,*” arXiv:0806.3253 [hep-th].
- [51] M. Blau and M. O’Loughlin, “*DLCQ and Plane Wave Matrix Big Bang Models,*” arXiv:0806.3255 [hep-th].
- [52] T. Nishioka and T. Takayanagi, “*On Type IIA Penrose Limit and N=6 Chern-Simons Theories,*” arXiv:0806.3391 [hep-th].
- [53] Y. Honma, S. Iso, Y. Sumitomo and S. Zhang, “*Scaling limit of N=6 superconformal Chern-Simons theories and Lorentzian Bagger-Lambert theories,*” arXiv:0806.3498 [hep-th].
- [54] Y. Imamura and K. Kimura, “*Coulomb branch of generalized ABJM models,*” arXiv:0806.3727 [hep-th].
- [55] J. A. Minahan and K. Zarembo, “*The Bethe ansatz for superconformal Chern-Simons,*” arXiv:0806.3951 [hep-th].
- [56] K. Furuuchi, S. Y. Shih and T. Takimi, “*M-Theory Superalgebra From Multiple Membranes,*” arXiv:0806.4044 [hep-th].
- [57] A. Agarwal, “*Mass Deformations of Super Yang-Mills Theories in D= 2+1, and Super-Membranes: A Note,*” arXiv:0806.4292 [hep-th].
- [58] D. Gaiotto, S. Giombi and X. Yin, “*Spin Chains in N=6 Superconformal Chern-Simons-Matter Theory,*” arXiv:0806.4589 [hep-th].
- [59] I. A. Bandos and P. K. Townsend, “*Light-cone M5 and multiple M2-branes,*” arXiv:0806.4777 [hep-th].

- [60] C. Ahn, “Towards Holographic Gravity Dual of $N=1$ Superconformal Chern-Simons Gauge Theory,” arXiv:0806.4807 [hep-th].
- [61] J. Bedford and D. Berman, “A note on Quantum Aspects of Multiple Membranes,” arXiv:0806.4900 [hep-th].
- [62] G. Arutyunov and S. Frolov, “Superstrings on $AdS_4 \times CP^3$ as a Coset Sigma-model,” arXiv:0806.4940 [hep-th].
- [63] B. . j. Stefanski, “Green-Schwarz action for Type IIA strings on $AdS_4 \times CP^3$,” arXiv:0806.4948 [hep-th].
- [64] G. Grignani, T. Harmark and M. Orselli, “The $SU(2) \times SU(2)$ sector in the string dual of $N=6$ superconformal Chern-Simons theory,” arXiv:0806.4959 [hep-th].
- [65] K. Hosomichi, K. M. Lee, S. Lee, S. Lee and J. Park, “ $N=5,6$ Superconformal Chern-Simons Theories and $M2$ -branes on Orbifolds,” arXiv:0806.4977 [hep-th].
- [66] P. Fre and P. A. Grassi, “Pure Spinor Formalism for $Osp(N-4)$ backgrounds,” arXiv:0807.0044 [hep-th].
- [67] J. Bagger and N. Lambert, “Three-Algebras and $N=6$ Chern-Simons Gauge Theories,” arXiv:0807.0163 [hep-th].
- [68] S. Terashima, “On $M5$ -branes in $N=6$ Membrane Action,” arXiv:0807.0197 [hep-th].
- [69] G. Grignani, T. Harmark, M. Orselli and G. W. Semenoff, “Finite size Giant Magnons in the string dual of $N=6$ superconformal Chern-Simons theory,” arXiv:0807.0205 [hep-th].
- [70] S. Terashima and F. Yagi, “Orbifolding the Membrane Action,” arXiv:0807.0368 [hep-th].
- [71] N. Gromov and P. Vieira, “The AdS_4/CFT_3 algebraic curve,” arXiv:0807.0437 [hep-th].
- [72] C. Ahn and P. Bozhilov, “Finite-size effects of Membranes on $AdS_4 \times S_7$,” arXiv:0807.0566 [hep-th].
- [73] N. Gromov and P. Vieira, “The all loop AdS_4/CFT_3 Bethe ansatz,” arXiv:0807.0777 [hep-th].
- [74] B. Chen and J. B. Wu, “Semi-classical strings in $AdS_4 * CP^3$,” arXiv:0807.0802 [hep-th].
- [75] S. Cherkis and C. Saemann, “Multiple $M2$ -branes and Generalized 3-Lie algebras,” arXiv:0807.0808 [hep-th].
- [76] C. S. Chu, P. M. Ho, Y. Matsuo and S. Shiba, “Truncated Nambu-Poisson Bracket and Entropy Formula for Multiple Membranes,” arXiv:0807.0812 [hep-th].
- [77] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, “Studies of the ABJM Theory in a Formulation with Manifest $SU(4)$ R-Symmetry,” arXiv:0807.0880 [hep-th].
- [78] Y. Zhou, “Interpreting N $M2$ branes,” arXiv:0807.0890 [hep-th].
- [79] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, “A Massive Study of $M2$ -brane Proposals,” arXiv:0807.1074 [hep-th].
- [80] M. Schnabl and Y. Tachikawa, “Classification of $N=6$ superconformal theories of ABJM type,” arXiv:0807.1102 [hep-th].
- [81] T. Li, Y. Liu and D. Xie, “Multiple $D2$ -Brane Action from $M2$ -Branes,” arXiv:0807.1183 [hep-th].

- [82] N. Kim, “How to put the Bagger-Lambert theory on an orbifold : A derivation of the ABJM model,” arXiv:0807.1349 [hep-th].
- [83] Y. Pang and T. Wang, “From N M2’s to N D2’s,” arXiv:0807.1444 [hep-th].
- [84] M. R. Garousi, A. Ghodsi and M. Khosravi, “On thermodynamics of $N=6$ superconformal Chern-Simons theories at strong coupling,” arXiv:0807.1478 [hep-th].
- [85] A. Hashimoto and P. Ouyang, “Supergravity dual of Chern-Simons Yang-Mills theory with $N=6,8$ superconformal IR fixed point,” arXiv:0807.1500 [hep-th].
- [86] D. Astolfi, V. G. M. Puletti, G. Grignani, T. Harmark and M. Orselli, “Finite-size corrections in the $SU(2) \times SU(2)$ sector of type IIA string theory on $AdS_4 \times CP^3$,” arXiv:0807.1527 [hep-th].
- [87] M. Ali-Akbari, M. M. Sheikh-Jabbari and J. Simon, “Relaxed Three-Algebras: Their Matrix Representations and Implications for Multi M2-brane Theory,” arXiv:0807.1570 [hep-th].
- [88] C. Ahn and R. I. Nepomechie, “ $N=6$ super Chern-Simons theory S -matrix and all-loop Bethe ansatz equations,” arXiv:0807.1924 [hep-th].
- [89] K. Hanaki and H. Lin, “M2-M5 Systems in $N=6$ Chern-Simons Theory,” arXiv:0807.2074 [hep-th].
- [90] H. Verlinde, “D2 or M2? A Note on Membrane Scattering,” arXiv:0807.2121 [hep-th].
- [91] Y. Imamura and K. Kimura, “ $N=4$ Chern-Simons theories with auxiliary vector multiplets,” arXiv:0807.2144 [hep-th].
- [92] Y. Lozano, “Eleven dimensions from the massive D-2-brane,” Phys. Lett. B **414** (1997) 52 [arXiv:hep-th/9707011].
- [93] T. Ortin, “A note on the D-2-brane of the massive type IIA theory and gauged sigma models,” Phys. Lett. B **415** (1997) 39 [arXiv:hep-th/9707113].
- [94] L. J. Romans, “Massive $N=2a$ Supergravity In Ten-Dimensions,” Phys. Lett. B **169** (1986) 374.
- [95] E. Bergshoeff, M. de Roo, M. B. Green, G. Papadopoulos and P. K. Townsend, “Duality of Type II 7-branes and 8-branes,” Nucl. Phys. B **470** (1996) 113 [arXiv:hep-th/9601150].
- [96] E. Eyras, B. Janssen and Y. Lozano, “5-branes, KK-monopoles and T-duality,” Nucl. Phys. B **531** (1998) 275 [arXiv:hep-th/9806169].
- [97] E. Bergshoeff and J. P. van der Schaar, “On M-9-branes,” Class. Quant. Grav. **16** (1999) 23 [arXiv:hep-th/9806069].
- [98] E. Bergshoeff, Y. Lozano and T. Ortin, “Massive branes,” Nucl. Phys. B **518** (1998) 363 [arXiv:hep-th/9712115].
- [99] E. Bergshoeff, P. M. Cowdall and P. K. Townsend, “Massive IIA supergravity from the topologically massive D-2-brane,” Phys. Lett. B **410** (1997) 13 [arXiv:hep-th/9706094].
- [100] Y. Honma, S. Iso, Y. Sumitomo, H. Umetsu and S. Zhang, “Generalized Conformal Symmetry and Recovery of $SO(8)$ in Multiple M2 and D2 Branes,” arXiv:0807.3825 [hep-th].
- [101] E. A. Bergshoeff, O. Hohm, D. Roest, H. Samtleben and E. Sezgin, “The Superconformal Gaugings in Three Dimensions,” arXiv:0807.2841 [hep-th].

- [102] B. H. Lee, K. L. Panigrahi and C. Park, “*Spiky Strings on $AdS_4 \times \mathbf{CP}^3$* ,” arXiv:0807.2559 [hep-th].
- [103] C. Krishnan and C. Maccaferri, “*Membranes on Calibrations*,” arXiv:0805.3125 [hep-th].